

Statistics for Business

Sampling Distributions, Interval Estimation and Hypothesis Tests.

Panagiotis Th. Konstantinou

MSc in International Shipping, Finance and Management,

Athens University of Economics and Business

First Draft: July 15, 2015. **This Draft:** August 28, 2023.

Lecture Outline

- Simple random sampling
- Distribution of the sample average
- Large sample approximation to the distribution of the sample mean
 - ▶ Law of Large Numbers
 - ▶ Central Limit Theorem
- Estimation of the population mean
 - ▶ Unbiasedness
 - ▶ Consistency
 - ▶ Efficiency
- Hypothesis test concerning the population mean
- Confidence intervals for the population mean
 - ▶ Using the t -statistic when n is small
- Comparing means from different populations

Sampling

- A **population** is a collection of all the elements of interest, while a **sample** is a subset of the population.
- The reason we select a sample is to collect data to answer a research question about a population.
- The sample results provide only **estimates** of the values of the population characteristics. With *proper sampling methods*, the sample results can provide “good” estimates of the population characteristics.
- A **random sample** from an infinite population is a sample selected such that the following conditions are satisfied:
 - ▶ Each element selected comes from the population of interest.
 - ▶ Each element is selected independently.
 - ★ If the population is finite, then we sample with replacement...

Simple Random Sampling – I

- **Simple random sampling** means that n objects are drawn randomly from a population and each object is equally likely to be drawn
- Let Y_1, Y_2, \dots, Y_n denote the 1st to the n th randomly drawn object. Under simple random sampling
 - ▶ The marginal probability distribution of Y_i is the same for all $i = 1, 2, \dots, n$ and equals the population distribution of Y .
 - ★ because Y_1, Y_2, \dots, Y_n are drawn randomly from the **same** population.
 - ▶ Y_1 is distributed independently from Y_2, \dots, Y_n . knowing the value of Y_i does not provide information on Y_j for $i \neq j$
- When Y_1, Y_2, \dots, Y_n are drawn from the same population and are independently distributed, they are said to be **I.I.D. random variables**

Simple Random Sampling – II

Example

- Let G be the gender of an individual ($G = 1$ if female, $G = 0$ if male)
- G is a Bernoulli r.v. with $E(G) = \mu_G = \Pr(G = 1) = 0.5$
- Suppose we take the population register and randomly draw a sample of size n
 - ▶ The probability distribution of G_i is a Bernoulli with mean 0.5
 - ▶ G_1 is distributed independently from G_2, \dots, G_n
- Suppose we draw a random sample of individuals entering the building of the accounting department
 - ▶ This is not a sample obtained by simple random sampling and G_1, G_2, \dots, G_n are not i.i.d
 - ▶ Men are more likely to enter the building of the accounting department!

The Sampling Distribution of the Sample Average – I

- The *sample average* \bar{Y} of a randomly drawn sample is a random variable with a probability distribution called the *sampling distribution*

$$\bar{Y} = \frac{1}{n}(Y_1 + Y_2 + \cdots + Y_n) = \frac{1}{n} \sum_{i=1}^n Y_i$$

- ▶ The individuals in the sample are drawn at random.
- ▶ Thus the values of (Y_1, Y_2, \cdots, Y_n) are random
- ▶ Thus functions of (Y_1, Y_2, \cdots, Y_n) , such as \bar{Y} , are random: had a different sample been drawn, they would have taken on a different value
- ▶ The distribution of over different possible samples of size n is called the *sampling distribution* of \bar{Y} .
- ▶ The mean and variance of are the mean and variance of its sampling distribution, $E(\bar{Y})$ and $\text{Var}(\bar{Y})$.
- ▶ The concept of the sampling distribution underpins all of statistics/econometrics.

The Sampling Distribution of the Sample Average – II

$$\bar{Y} = \frac{1}{n}(Y_1 + Y_2 + \cdots + Y_n) = \frac{1}{n} \sum_{i=1}^n Y_i$$

- Suppose that Y_1, Y_2, \dots, Y_n are *I.I.D.* and the mean & variance of the population distribution of Y are respectively μ_Y and σ_Y^2

- ▶ The mean of (the sampling distribution of) \bar{Y} is

$$E(\bar{Y}) = E\left(\frac{1}{n} \sum_{i=1}^n Y_i\right) = \frac{1}{n} \sum_{i=1}^n E(Y_i) = \frac{1}{n} n E(Y) = \mu_Y$$

- ▶ The variance of (the sampling distribution of) \bar{Y} is

$$\begin{aligned} \text{Var}(\bar{Y}) &= \text{Var}\left(\frac{1}{n} \sum_{i=1}^n Y_i\right) = \frac{1}{n^2} \sum_{i=1}^n \text{Var}(Y_i) + 2 \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1, j \neq i}^n \text{Cov}(Y_i, Y_j) \\ &= \frac{1}{n^2} n \text{Var}(Y) + 0 = \frac{1}{n} \text{Var}(Y) = \frac{\sigma_Y^2}{n} \end{aligned}$$

The Sampling Distribution of the Sample Average – III

Example

- Let G be the gender of an individual ($G = 1$ if female, $G = 0$ if male)
- The mean of the population distribution of G is

$$E(G) = \mu_G = \Pr(G = 1) = p = 0.5$$

- The variance of the population distribution of G is

$$\text{Var}(G) = \sigma_G^2 = p(1 - p) = 0.5(1 - 0.5) = 0.25$$

- The mean and variance of the average gender (proportion of women) \bar{G} in a random sample with $n = 10$ are

$$\begin{aligned} E(\bar{G}) &= \mu_G = 0.5 \\ \text{Var}(\bar{G}) &= \frac{1}{n} \sigma_G^2 = \frac{1}{10} 0.25 = 0.025 \end{aligned}$$

The Finite-Sample Distribution of the Sample Average

- The *finite sample distribution* is the sampling distribution that exactly describes the distribution of \bar{Y} for any sample size n .
- In general the exact sampling distribution of \bar{Y} is complicated and depends on the population distribution of Y .
- A special case is when Y_1, Y_2, \dots, Y_n are *IID* draws from the $N(\mu_Y, \sigma_Y^2)$, because in this case

$$\bar{Y} \sim N\left(\mu_Y, \frac{\sigma_Y^2}{n}\right)$$

The Sampling Distribution of the Average Gender \bar{G}

- Suppose G takes on 0 or 1 (a Bernoulli random variable) with the probability distribution

$$\Pr(G = 0) = p = 0.5, \quad \Pr(G = 1) = 1 - p = 0.5$$

- As we discussed above:


$$\begin{aligned} E(G) &= \mu_G = \Pr(G = 1) = p = 0.5 \\ \text{Var}(G) &= \sigma_G^2 = p(1 - p) = 0.5(1 - 0.5) = 0.25 \end{aligned}$$

- The sampling distribution of \bar{G} depends on n .
- Consider $n = 2$. The sampling distribution of \bar{G} is
 - ▶ $\Pr(\bar{G} = 0) = 0.5^2 = 0.25$
 - ▶ $\Pr(\bar{G} = 1/2) = 2 \times 0.5 \times (1 - 0.5) = 0.5$
 - ▶ $\Pr(\bar{G} = 1) = (1 - 0.5)^2 = 0.25$

The Finite-Sample Distribution of the Average Gender \bar{G}

- Suppose we draw 999 samples of $n = 2$:

Sample 1			Sample 1			Sample 3			...	Sample 999		
G_1	G_2	\bar{G}	G_1	G_2	\bar{G}	G_1	G_2	\bar{G}		G_1	G_2	\bar{G}
1	0	0.5	1	1	1	0	1	0.5		0	0	0



The Asymptotic Distribution of the Sample Average \bar{Y}

- Given that the exact sampling distribution of \bar{Y} is complicated and given that we generally use large samples in statistics/econometrics we will often use an approximation of the sample distribution that relies on the sample being large
- The *asymptotic distribution* or *large-sample distribution* is the approximate sampling distribution of \bar{Y} if the sample size becomes very large: $n \rightarrow \infty$.
- We will use two concepts to approximate the large-sample distribution of the sample average
 - ▶ The law of large numbers.
 - ▶ The central limit theorem.

The Law of Large Numbers (LLN)

Definition (Law of Large Numbers)

Suppose that

- 1 $Y_i, i = 1, \dots, n$ are independently and identically distributed with $E(Y_i) = \mu_Y$; and
- 2 large outliers are unlikely i.e. $\text{Var}(Y_i) = \sigma_Y^2 < +\infty$.

Then \bar{Y} will be near μ_Y with very high probability when n is very large ($n \rightarrow \infty$)

$$\bar{Y} \xrightarrow{p} \mu_Y.$$





We also say that the sequence of random variables $\{Y_n\}$ converges in probability to the μ_Y , if for every $\varepsilon > 0$

$$\lim_{n \rightarrow \infty} \Pr(|\bar{Y}_n - \mu_Y| > \varepsilon) = 0.$$

We also denote this by $\text{plim}(Y_n) = \mu_Y$

The Law of Large Numbers (LLN)

Example: Gender $G \sim \text{Bernoulli}(0.5, 0.25)$



The Central Limit Theorem (CLT)

Definition (Central Limit Theorem)

Suppose that

- 1 $Y_i, i = 1, \dots, n$ are independently and identically distributed with $E(Y_i) = \mu_Y$; and
- 2 large outliers are unlikely i.e. $\text{Var}(Y_i) = \sigma_Y^2$ with $0 < \sigma_Y^2 < +\infty$.

Then the distribution of the sample average \bar{Y} will be approximately normal as n becomes very large ($n \rightarrow \infty$)





$$\bar{Y} \sim N\left(\mu_Y, \frac{\sigma_Y^2}{n}\right).$$

The distribution of the the standardized sample average is approximately standard normal for $n \rightarrow \infty$

$$\frac{\bar{Y} - \mu_Y}{\sigma_Y/\sqrt{n}}$$

The Central Limit Theorem (CLT)

Example: Gender $G \sim \text{Bernoulli}(0.5, 0.25)$



The Central Limit Theorem (CLT)

- How good is the large-sample approximation?
- ★ If $Y_i \sim N(\mu_Y, \sigma_Y^2)$ the approximation is perfect.
- ★ If Y_i is not normally distributed the quality of the approximation depends on how close n is to infinity (how large n is)
- ★ For $n \geq 100$ the normal approximation to the distribution of \bar{Y} is typically very good for a wide variety of population distributions.

Estimators and Estimates

Definition

An **estimator** is a function of a sample of data to be drawn randomly from a population.

- An estimator is a random variable because of randomness in drawing the sample. Typically used estimators

$$\text{Sample Average: } \bar{Y} = \frac{1}{n} \sum_{i=1}^n Y_i, \text{ Sample variance: } S_Y^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i - \bar{Y})^2.$$

Using a particular sample y_1, y_2, \dots, y_n we obtain

$$\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i \text{ and } s_y^2 = \frac{1}{n-1} \sum_{i=1}^n (y_i - \bar{y})^2$$

which are **point estimates**. These are the numerical value of an estimator when it is actually computed using a specific sample.

Estimation of the Population Mean – I

- Suppose we want to know the mean value of Y (μ_Y) in a population, for example
 - ▶ The mean wage of college graduates.
 - ▶ The mean level of education in Greece.
 - ▶ The mean probability of passing the statistics exam.
- Suppose we draw a random sample of size n with Y_1, Y_2, \dots, Y_n being *IID*
- Possible estimators of μ_Y are:
 - ▶ The sample average: $\bar{Y} = \frac{1}{n} \sum_{i=1}^n Y_i$
 - ▶ The first observation: Y_1
 - ▶ The weighted average: $\tilde{Y} = \frac{1}{n} \left(\frac{1}{2} Y_1 + \frac{3}{2} Y_2 + \dots + \frac{1}{2} Y_{n-1} + \frac{3}{2} Y_n \right)$.
- To determine which of the estimators, \bar{Y} , Y_1 or \tilde{Y} is the best estimator of μ_Y we consider 3 properties.
- Let $\hat{\mu}_Y$ be an estimator of the population mean μ_Y

Estimation of the Population Mean – II

- ① **Unbiasedness:** The mean of the sampling distribution of $\hat{\mu}_Y$ equals μ_Y

$$E(\hat{\mu}_Y) = \mu_Y.$$

- ② **Consistency:** The probability that $\hat{\mu}_Y$ is within a very small interval of μ_Y approaches 1 if $n \rightarrow \infty$

$$\hat{\mu}_Y \xrightarrow{P} \mu_Y \text{ or } \Pr(|\hat{\mu}_Y - \mu_Y| < \varepsilon) = 1$$

- ③ **Efficiency:** If the variance of the sampling distribution of $\hat{\mu}_Y$ is smaller than that of some other estimator $\tilde{\mu}_Y$, $\hat{\mu}_Y$ is more efficient

$$\text{Var}(\hat{\mu}_Y) \leq \text{Var}(\tilde{\mu}_Y)$$

Estimating Mean Wages – I

- Suppose we are interested in the mean wages (pre tax) μ_W of individuals with a Ph.D. in economics/finance in Europe (true mean $\mu_w = 60K$). We draw the following sample ($n = 10$) by simple random sampling

i	1	2	3	4	5
W_i	47281.92	70781.94	55174.46	49096.05	67424.82
i	6	7	8	9	10
W_i	39252.85	78815.33	46750.78	46587.89	25015.71

- The 3 estimators give the following estimates:

- ▶ $\bar{W} = \frac{1}{10} \sum_{i=1}^{10} W_i = 52618.18$

- ▶ $W_1 = 47281.92$



- ▶ $\tilde{W} = \frac{1}{10} \left(\frac{1}{2} W_1 + \frac{3}{2} W_2 + \dots + \frac{1}{2} W_9 + \frac{3}{2} W_{10} \right) = 49398.82$

- Unbiasedness:** All 3 proposed estimators are unbiased

Estimating Mean Wages – II

- Consistency:**


- ▶ By the law of large numbers $\bar{W} \xrightarrow{P} \mu_W$ which implies that the probability that \bar{W} is within a very small interval of μ_W approaches 1 if $n \rightarrow \infty$




Estimating Mean Wages – III

- ▶ $\tilde{W} = \frac{1}{n} \left(\frac{1}{2} W_1 + \frac{3}{2} W_2 + \dots + \frac{1}{2} W_{n-1} + \frac{3}{2} W_n \right)$ can also be shown to be consistent

Weighted average as estimator of population mean
999 samples of n=10




Weighted average as estimator of population mean
999 samples of n=100




- ▶ However W_1 is not a consistent estimator of μ_W .

First observation W_1 as estimator of population mean
999 samples of n=10



First observation W_1 as estimator of population mean
999 samples of n=100



Estimating Mean Wages – IV

- **Efficiency:** We have that
 - ▶ $\text{Var}(\bar{W}) = \frac{1}{n}\sigma_W^2$
 - ▶ $\text{Var}(W_1) = \sigma_W^2$
 - ▶ $\text{Var}(\tilde{W}) = 1.25\frac{1}{n}\sigma_W^2$
 - ▶ So for any $n \geq 2$, \bar{W} is more efficient than W_1 and \tilde{W} .
- In fact \bar{Y} is the **Best Linear Unbiased Estimator (BLUE)**: it is the most efficient estimator of μ_Y among all unbiased estimators that are weighted averages of Y_1, Y_2, \dots, Y_n
- ★ Let $\hat{\mu}_Y = \frac{1}{n} \sum_{i=1}^n \alpha_i Y_i$ be an unbiased estimator of μ_Y with α_i nonrandom constants. Then \bar{Y} is more efficient than $\hat{\mu}_Y$

$$\text{Var}(\bar{Y}) \leq \text{Var}(\hat{\mu}_Y)$$

Hypothesis Tests

Consider the following questions:

- Is the mean monthly wage of Ph.D. graduates equal to 60000 euros?
- Is the mean level of education in Greece equal to 12 years?
- Is the mean probability of passing the stats exam equal to 1?

These questions involve the population mean taking on a specific value $\mu_{Y,0}$. Answering these questions implies using data to compare a **null hypothesis** (a tentative assumption about the population mean parameter)

$$H_0 : E(Y) = \mu_{Y,0}$$

to an **alternative hypothesis** (the opposite of what is stated in the H_0)

$$H_1 : E(Y) \neq \mu_{Y,0}$$

- Alternative Hypothesis as a Research Hypothesis

- ▶ **Example:** A new sales force bonus plan is developed in an attempt to increase sales.
- ▶ **Alternative Hypothesis:** The new bonus plan increase sales.
- ▶ **Null Hypothesis:** The new bonus plan does not increase sales.

Hypothesis Tests: Terminology

- The **hypothesis testing problem** (for the mean): make a provisional decision, based on the evidence at hand, whether a null hypothesis is true, or instead that some alternative hypothesis is true. That is, test
 - ▶ $H_0 : E(Y) \leq \mu_{Y,0}$ vs. $H_1 : E(Y) > \mu_{Y,0}$ (1-sided, $>$)
 - ▶ $H_0 : E(Y) \geq \mu_{Y,0}$ vs. $H_1 : E(Y) < \mu_{Y,0}$ (1-sided, $<$)
 - ▶ $H_0 : E(Y) = \mu_{Y,0}$ vs. $H_1 : E(Y) \neq \mu_{Y,0}$ (2-sided)
- p -value = probability of drawing a statistic (e.g. \bar{Y}) at least as adverse to the null as the value actually computed with your data, assuming that the null hypothesis is true.
- The **significance level** of a test (α) is a pre-specified probability of incorrectly rejecting the null, when the null is true. Typical values are 0.01 (1%), 0.05 (5%), or 0.10 (10%).
 - ▶ It is selected by the researcher at the beginning, and determines the **critical value(s)** of the test.
 - ▶ If the test-statistic falls outside the non-rejection region, we reject H_0 .

Hypothesis Tests

The Testing Process and Rejections


Level of significance = α

★ Represents critical value

$$H_0: E(Y) = \mu_{Y,0}$$

$$H_1: E(Y) \neq \mu_{Y,0}$$

Two-tail test




Rejection region is shaded

$$H_0: E(Y) \leq \mu_{Y,0}$$

$$H_1: E(Y) > \mu_{Y,0}$$


Right-tail test



$$H_0: E(Y) \geq \mu_{Y,0}$$

$$H_1: E(Y) < \mu_{Y,0}$$

Left-tail test



Hypothesis Testing using p -values

- The p -value is the probability, computed using the test statistic, that measures the support (or lack of support) provided by the sample for the null hypothesis
 - ▶ If the p -value is less than or equal to the level of significance α , the value of the test statistic is in the rejection region.
 - ▶ Reject H_0 if the p -value $< \alpha$.
 - ▶ See also Annex
- **Rules of thumb**
 - ▶ If p -value is less than .01, there is overwhelming evidence to conclude H_0 is false.
 - ▶ If p -value is between .01 and .05, there is strong evidence to conclude H_0 is false.
 - ▶ If p -value is between .05 and .10, there is weak evidence to conclude H_0 is false.
 - ▶ If p -value is greater than .10, there is insufficient evidence to conclude H_0 is false.

Hypothesis Test for the Mean with σ_Y^2 known – I

Decision Rules

- The test statistic employed is obtained by converting the sample result (\bar{y}) to a z -value

$$z = \frac{\bar{y} - \mu_{Y,0}}{\sigma_Y / \sqrt{n}}$$

$$\begin{array}{l} H_0 : E(Y) \geq \mu_{Y,0} \\ H_1 : E(Y) < \mu_{Y,0} \end{array}$$

Lower-tail

Reject H_0 if $z < z_\alpha$

$$\begin{array}{l} H_0 : E(Y) \leq \mu_{Y,0} \\ H_1 : E(Y) > \mu_{Y,0} \end{array}$$

Upper-tail

Reject H_0 if $z > z_\alpha$

$$\begin{array}{l} H_0 : E(Y) = \mu_{Y,0} \\ H_1 : E(Y) \neq \mu_{Y,0} \end{array}$$

Two-tailed

Reject H_0 if $z < -z_{\alpha/2}$
or if $z > z_{\alpha/2}$

Hypothesis Test for the Mean with σ_Y^2 known – II


Decision Rules

$$\text{Hypothesis Tests for } E(Y) \quad z = \frac{\bar{Y} - \mu_{Y,0}}{\sigma_{\bar{Y}}} = \frac{\bar{Y} - \mu_{Y,0}}{\sigma_Y/\sqrt{n}}$$

Lower-tail test:

$$H_0: E(Y) \geq \mu_0$$

$$H_1: E(Y) < \mu_0$$




Reject H_0 if $z < -z_\alpha$

Upper-tail test:

$$H_0: E(Y) \leq \mu_{Y,0}$$

$$H_1: E(Y) > \mu_{Y,0}$$




Reject H_0 if $z > z_\alpha$

Two-tail test:

$$H_0: E(Y) = \mu_{Y,0}$$

$$H_1: E(Y) \neq \mu_{Y,0}$$



Reject H_0 if $z < -z_{\alpha/2}$
or $z > z_{\alpha/2}$

Hypothesis Test for the Mean (σ^2 known) – I

Examples

- Example 1.** A phone industry manager thinks that customer monthly cell phone bill have increased, and now average over \$52 per month. The company wishes to test this claim. Assume $\sigma = 10$ is known and let $\alpha = 0.10$. Suppose a sample of 64 persons is taken, and it is found that the average bill \$53.1.

- ▶ Form the hypothesis to be tested

$$H_0 : E(Y) \leq 52 \quad \text{the mean is not over } \$52 \text{ per month}$$

$$H_1 : E(Y) > 52 \quad \text{the mean is over } \$52 \text{ per month}$$

- ▶ For $\alpha = 0.10$, $z_{0.10} = 1.28$, so we would reject H_0 if $z > 1.28$.
- ▶ We have $n = 64$ and $\bar{y} = 53.1$, so the test statistic is

$$z = \frac{\bar{y} - \mu_{Y,0}}{\sigma_Y / \sqrt{n}} = \frac{53.1 - 52}{10 / \sqrt{64}} = 0.88 < z_{0.10} = 1.28$$

Hence H_0 cannot be rejected.

Hypothesis Test for the Mean (σ^2 known) – II

Examples

- **Example 2.** We would like to test the claim that the true mean # of TV sets in EU homes is equal to 3 (assuming $\sigma_Y = 0.8$ known). For this purpose a sample of 100 homes is selected, and the average number of TV sets is 2.84. Test the above hypothesis using $\alpha = 0.05$.

- ▶ Form the hypothesis to be tested

$H_0 : E(Y) = 3$ the *mean # is* 3 TV sets per home

$H_1 : E(Y) \neq 3$ the *mean is not* 3 TV sets per home

- ▶ For $\alpha = 0.05$, $z_{\alpha/2} = z_{0.025} = 1.96$ and $-z_{0.025} = -1.96$, so we would reject H_0 if $|z| > 1.96$.

Hypothesis Test for the Mean (σ^2 known) – III

Examples

- ▶ We have $n = 100$ and $\bar{y} = 2.84$, so the test statistic is

$$z = \frac{\bar{y} - \mu_{Y,0}}{\sigma_Y / \sqrt{n}} = \frac{2.84 - 3}{0.8 / \sqrt{100}} = \frac{-0.16}{0.08} = -2 < -z_{0.025} = -1.96$$

or $|z| = 2 > 1.96$, Hence H_0 is rejected. We **conclude** that there is sufficient evidence that the mean number of TVs in EU homes is not equal to 3.

Test for the Mean with σ_Y^2 unknown but $n \rightarrow \infty$

Decision Rules


- Since $S_Y^2 \xrightarrow{P} \sigma_Y^2$, compute the standard error of \bar{Y} , $SE(\bar{Y}) = s_Y/\sqrt{n}$ and construct a t -ratio.

$$\text{Hypothesis Tests for } E(Y) \quad t = \frac{\bar{Y} - \mu_{Y,0}}{SE(\bar{Y})} = \frac{\bar{Y} - \mu_{Y,0}}{s_Y/\sqrt{n}}$$

Lower-tail test:

$$H_0: E(Y) \geq \mu_0$$

$$H_1: E(Y) < \mu_0$$




$$-z_\alpha$$

Reject H_0 if $t < -z_\alpha$

Upper-tail test:

$$H_0: E(Y) \leq \mu_{Y,0}$$

$$H_1: E(Y) > \mu_{Y,0}$$




$$z_\alpha$$

Reject H_0 if $t > z_\alpha$

Two-tail test:

$$H_0: E(Y) = \mu_{Y,0}$$

$$H_1: E(Y) \neq \mu_{Y,0}$$



$$-z_{\alpha/2}$$

$$z_{\alpha/2}$$

Reject H_0 if $t < -z_{\alpha/2}$
or $t > z_{\alpha/2}$

Test for the Mean with σ_Y^2 unknown but $n \rightarrow \infty$

Example

- Suppose we would like to test

$$H_0 : E(W) = 60000, \quad H_1 : E(W) \neq 60000,$$

using a sample of 250 individuals with a Ph.D. degree at the 5% significance level.

- We perform the following steps:

- $\bar{W} = \frac{1}{n} \sum_{i=1}^n W_i = \frac{1}{250} \sum_{i=1}^{250} W_i = 61977.12.$

- $SE(\bar{W}) = \frac{s_W}{\sqrt{n}} = \frac{s_W}{\sqrt{250}} = 1334.19.$

- Compute $t^{act} = \frac{\bar{W} - \mu_{W,0}}{SE(\bar{W})} = \frac{61977.12 - 60000}{1334.19} = 1.4819.$

- Since we use a 5% significance level, we do not reject H_0 because $|t^{act}| = 1.4819 < z_{0.025} = 1.96.$

- Suppose we are interested in the alternative $H_1 : E(W) > 60000$. The t -stat is **exactly** the same: $t^{act} = 1.4819$. but now needs to be compared with $z_{0.05} = 1.645$.

Hypothesis Test for the Mean with σ^2 unknown (n small)

Decision Rules


- Consider a random sample of n observations from a population that is normally distributed, **AND** variance σ_Y^2 is unknown: $Y_i \sim N(\mu_Y, \sigma_Y^2)$
- Converting the sample average (\bar{y}) to a t -value...

$$\text{Hypothesis Tests for } E(Y) \quad t = \frac{\bar{Y} - \mu_{Y,0}}{\text{SE}(\bar{Y})} = \frac{\bar{Y} - \mu_{Y,0}}{s_Y/\sqrt{n}} \sim t_{n-1}$$

Lower-tail test:

$$H_0: E(Y) \geq \mu_0$$

$$H_1: E(Y) < \mu_0$$




Reject H_0 if $t < -t_{n-1, \alpha}$

Upper-tail test:

$$H_0: E(Y) \leq \mu_0$$

$$H_1: E(Y) > \mu_0$$




Reject H_0 if $t > t_{n-1, \alpha}$

Two-tail test:

$$H_0: E(Y) = \mu_0$$

$$H_1: E(Y) \neq \mu_0$$



Reject H_0 if $t < -t_{n-1, \alpha/2}$
or $t > t_{n-1, \alpha/2}$

Hypothesis Test for the Mean with σ^2 unknown (n small)

Example

- The average cost of a hotel room in New York is said to be \$168 per night. A random sample of 25 hotels resulted in $\bar{y} = \$172.50$ and $s_y = \$15.40$. Perform a test at the $\alpha = 0.05$ level (assuming the population distribution is normal).
 - Form the hypothesis to be tested

$$H_0 : E(Y) = 168 \quad \text{the mean cost is \$168}$$

$$H_1 : E(Y) \neq 168 \quad \text{the mean cost is not \$168}$$

- For $\alpha = 0.05$, with $n = 25$, $t_{n-1, \alpha/2} = t_{24, 0.025} = 2.0639$ and $-t_{24, 0.025} = -2.0639$, so we would reject H_0 if $|t| > 2.0639$.
- We have $\bar{y} = 172.50$ and $s_y = 15.40$, so the test statistic is

$$t = \frac{\bar{y} - \mu_{Y,0}}{s_y / \sqrt{n}} = \frac{172.50 - 168}{15.40 / \sqrt{25}} = 1.46 < t_{24, 0.025} = 2.0639$$

or $|t| = 1.46 < 2.0639$. Hence H_0 **cannot be rejected**. We **conclude** that there is not sufficient evidence that the true mean cost is different than \$168.

Confidence Intervals for the Population Mean – I

- Suppose we would do a two-sided hypothesis test for many different values of $\mu_{0,Y}$. On the basis of this we can construct a set of values which are not rejected at 5% ($\alpha\%$) significance level.
- If we were able to test all possible values of $\mu_{0,Y}$ we could construct a 95% ($(1 - \alpha)\%$) confidence interval

Definition

A 95% ($(1 - \alpha)\%$) confidence interval is an interval that contains the true value of μ_Y in 95% ($(1 - \alpha)\%$) of all possible random samples.

- ▶ A relative frequency interpretation: From repeated samples, 95% of all the confidence intervals that can be constructed will contain the unknown true population mean

Confidence Intervals for the Population Mean – II

- The general formula for all confidence intervals is

$$\text{Point Estimate} \pm \underbrace{(\text{Reliability Factor})(\text{Standard Error})}_{\text{Margin of Error}}$$

$$\hat{\mu} \pm c \cdot \text{SE}(\hat{\mu})$$

and using the sample average estimator

$$\bar{Y} \pm c \cdot \text{SE}(\bar{Y})$$

- Instead of doing infinitely many hypothesis tests we can compute the 95% $((1 - \alpha)\%)$ confidence interval as

$$\bar{Y} - z_{\alpha/2}\text{SE}(\bar{Y}) < \mu < \bar{Y} + z_{\alpha/2}\text{SE}(\bar{Y}) \quad \text{or} \quad \bar{Y} \pm \underbrace{z_{\alpha/2}\text{SE}(\bar{Y})}_{\text{Margin of Error}}$$

Confidence Intervals for the Population Mean – III

- When the sample size n is large (or when the population is normal and σ_Y^2 is known):
 - ▶ A 90% confidence interval for μ_Y : $[\bar{Y} \pm 1.645 \cdot \text{SE}(\bar{Y})]$
 - ▶ A 95% confidence interval for μ_Y : $[\bar{Y} \pm 1.96 \cdot \text{SE}(\bar{Y})]$
 - ▶ A 99% confidence interval for μ_Y : $[\bar{Y} \pm 2.58 \cdot \text{SE}(\bar{Y})]$

 - ▶ with $\text{SE}(\bar{Y}) = \sigma_Y/\sqrt{n}$ when variance is known or $\text{SE}(\bar{Y}) = s_Y/\sqrt{n}$ when unknown and is estimated.

Confidence Intervals for the Population Mean – IV

Example

A sample of 11 circuits from a large normal population has a mean resistance of 2.20 ohms. We know from past testing that the population standard deviation is 0.35 ohms. Determine a 95% C.I. for the true mean resistance of the population.

$$\begin{aligned}\bar{y} \pm z_{\alpha/2} \frac{\sigma_Y}{\sqrt{n}} &= 2.20 \pm 1.96(0.35/\sqrt{11}) = 2.20 \pm 0.2068 \\ 1.9932 &< \mu_Y < 2.4068\end{aligned}$$

- ▶ We are 95% confident that the true mean resistance is between 1.9932 and 2.4068 ohms
- ▶ Although the true mean may or may not be in this interval, 95% of intervals formed in this manner will contain the true mean

Confidence Intervals for the Population Mean – V

Example

Using the sample of $n = 250$ individuals with a Ph.D. degree discussed above ($\bar{W} = 61977.12$, $s_W = 21095.37$, $SE(\bar{Y}) = s_W/\sqrt{n} = 21095.37/\sqrt{250}$):

- ▶ A 90% C.I. for μ_W is: $[61977.12 \pm 1.64 \cdot 1334.19] = [59349.39, 64604.85]$.
- ▶ A 95% C.I. for μ_W is: $[61977.12 \pm 1.96 \cdot 1334.19] = [59774.38, 64179.86]$.
- ▶ A 99% C.I. for μ_W is: $[61977.12 \pm 2.58 \cdot 1334.19] = [58513.94, 65440.30]$.

Confidence Intervals for the Population Mean – VI

- When the sample size n is small **AND** the population from which we draw data is normal:

$$\bar{Y} - t_{n-1, \alpha/2} \frac{s_Y}{\sqrt{n}} < \mu_Y < \bar{Y} + t_{n-1, \alpha/2} \frac{s_Y}{\sqrt{n}} \quad \text{or} \quad \bar{Y} \pm \underbrace{t_{n-1, \alpha/2} \frac{s_Y}{\sqrt{n}}}_{\text{Margin of Error}}$$

- ▶ A 90% confidence interval for μ_Y : $[\bar{Y} \pm t_{n-1, 0.05} \cdot \text{SE}(\bar{Y})]$
- ▶ A 95% confidence interval for μ_Y : $[\bar{Y} \pm t_{n-1, 0.025} \cdot \text{SE}(\bar{Y})]$
- ▶ A 99% confidence interval for μ_Y : $[\bar{Y} \pm t_{n-1, 0.005} \cdot \text{SE}(\bar{Y})]$
- ▶ with $\text{SE}(\bar{Y}) = s_Y / \sqrt{n}$

Confidence Intervals for the Population Mean – VII

Example

A random sample of $n = 25$ has $\bar{x} = 50$ and $s = 8$. Form a 95% confidence interval for μ .

► $d.f. = n - 1 = 24$, so $t_{24, \alpha/2} = t_{24, 0.025} = 2.0639$

$$\begin{aligned}\bar{x} \pm t_{n-1, \alpha/2} \frac{s}{\sqrt{n}} &= 50 \pm 2.0639(8/\sqrt{25}) = 50 \pm 3.302 \\ 46.698 &< \mu < 53.302\end{aligned}$$

Comparing Means from Different Populations – I

Large Samples or Known Variances from Normal Populations

- Suppose we would like to test whether the mean wages of men and women with a Ph.D. degree differ by an amount d_0 :

$$H_0 : \mu_{W,M} - \mu_{W,F} = d_0 \quad H_0 : \mu_{W,M} - \mu_{W,F} \neq d_0$$

- To test the null hypothesis against the two-sided alternative we follow the 4 steps as above with some adjustments
- ① Estimate $(\mu_{W,M} - \mu_{W,F})$ by $(\bar{W}_M - \bar{W}_F)$.
 - ▶ Because a weighted average of 2 independent normal random variables is itself normally distributed we have (using the CLT and the fact that $\text{Cov}(\bar{W}_M, \bar{W}_F) = 0$)

$$\bar{W}_M - \bar{W}_F \sim N \left(\mu_{W,M} - \mu_{W,F}, \frac{\sigma_{W,M}^2}{n_M} + \frac{\sigma_{W,F}^2}{n_F} \right)$$

Comparing Means from Different Populations – II

Large Samples or Known Variances from Normal Populations

- 2 Estimate $\sigma_{W,M}$ and $\sigma_{W,F}$ to obtain $SE(\bar{W}_M - \bar{W}_F)$:

$$SE(\bar{W}_M - \bar{W}_F) = \sqrt{\frac{s_{W,M}^2}{n_M} + \frac{s_{W,F}^2}{n_F}}$$

- 3 Compute the t -statistic

$$t^{act} = \frac{(\bar{W}_M - \bar{W}_F) - d_0}{SE(\bar{W}_M - \bar{W}_F)}$$

- 4 Reject H_0 at a 5% significance level if $|t^{act}| > 1.96$ or if the p -value < 0.05 .

Comparing Means from Different Populations – III

Large Samples or Known Variances from Normal Populations

Example

Suppose we have random samples of 500 men and 500 women with a Ph.D. degree and we would like to test that the mean wages are equal:

$$H_0 : \mu_{W,M} - \mu_{W,M} = 0 \quad H_1 : \mu_{W,M} - \mu_{W,M} \neq 0$$

We obtained $\bar{W}_M = 64159.45$, $\bar{W}_F = 53163.41$, $s_{W,M} = 18957.26$, and $s_{W,F} = 20255.89$. We have:

① $\bar{W}_M - \bar{W}_F = 64159.45 - 53163.41 = 10996.04$.

② $SE(\bar{W}_M - \bar{W}_F) = 1240.709$.

③ $t^{act} = \frac{(\bar{W}_M - \bar{W}_F) - 0}{SE(\bar{W}_M - \bar{W}_F)} = \frac{10996.04}{1240.709} = 8.86$.

④ Since we use a 5% significance level, we reject H_0 because $|t^{act}| = 8.86 > 1.96$

Confidence Interval for the Difference in Population Means

- The method for constructing a confidence interval for 1 population mean can be easily extended to the difference between 2 population means.
- A hypothesized value of the difference in means d_0 will be rejected if $|t| > 1.96$ and will be in the confidence set if $|t| \leq 1.96$.
- Thus the 95% confidence interval for $\mu_{W,M} - \mu_{W,F}$ are the values of d_0 within ± 1.96 standard errors of $(\bar{W}_M - \bar{W}_F)$.
- So a 95% confidence interval for $\mu_{W,M} - \mu_{W,F}$ is

$$\begin{aligned} & (\bar{W}_M - \bar{W}_F) \pm 1.96 \cdot \text{SE}(\bar{W}_M - \bar{W}_F) \\ & 10996.04 \pm 1.96 \cdot 1240.709 \\ & [8561.34, 13430.73] \end{aligned}$$

Testing Population Mean Differences

Normal Populations, **Unknown Variances** σ_X^2 and σ_Y^2 but Assumed **Equal**

$$t = \frac{(\bar{X} - \bar{Y}) - d_0}{\text{SE}(\bar{X} - \bar{Y})} = \frac{(\bar{X} - \bar{Y}) - d_0}{\sqrt{(s_p^2/n_X) + (s_p^2/n_Y)}} \sim t_{n_X+n_Y-2};$$

$$\text{where } s_p^2 = \frac{(n_X - 1)s_X^2 + (n_Y - 1)s_Y^2}{n_X + n_Y - 2}$$

- The C.I. is constructed as $(\bar{X} - \bar{Y}) \pm t_{n_X+n_Y-2, \alpha/2} \cdot \text{SE}(\bar{X} - \bar{Y})$.
- Recall $\mu_X = E(X)$, $\mu_Y = E(Y)$

$$\begin{array}{l} H_0 : \mu_X - \mu_Y \geq d_0 \\ H_1 : \mu_X - \mu_Y < d_0 \end{array}$$

Lower-tail

Reject H_0 if $t < t_\alpha$

$$\begin{array}{l} H_0 : \mu_X - \mu_Y \leq d_0 \\ H_1 : \mu_X - \mu_Y > d_0 \end{array}$$

Upper-tail

Reject H_0 if $t > t_\alpha$

$$\begin{array}{l} H_0 : \mu_X - \mu_Y = d_0 \\ H_1 : \mu_X - \mu_Y \neq d_0 \end{array}$$

Two-tailed

Reject H_0 if $|t| > t_{\alpha/2}$

Testing Population Mean Differences – I

Example: Normal Populations, **Unknown Variances** σ_X^2 and σ_Y^2 but Assumed **Equal**

- You are a financial analyst for a brokerage firm. Is there a difference in dividend yield between stocks listed on the NYSE & NASDAQ? You collect the following data:

	NYSE	NASDAQ
Number:	21	25
Sample mean:	3.27	2.53
Sample std. dev.:	1.30	1.16

Assuming both populations are approximately normal with equal variances, is there a difference in average yield ($\alpha = 0.05$)?

- ▶ The hypothesis of interest is

$$\begin{array}{l} H_0 : \mu_{NYSE} - \mu_{NASDAQ} = 0 \\ H_1 : \mu_{NYSE} - \mu_{NASDAQ} \neq 0 \end{array} \text{ or } \begin{array}{l} H_0 : \mu_{NYSE} = \mu_{NASDAQ} \\ H_1 : \mu_{NYSE} \neq \mu_{NASDAQ} \end{array}$$

Testing Population Mean Differences – II

Example: Normal Populations, **Unknown Variances** σ_X^2 and σ_Y^2 but Assumed Equal

- ▶ Note that $df = n_X + n_Y - 2 = 21 + 25 - 2 = 44$, so the critical value for the test is $t_{44,0.025} = 2.0154$
- ▶ The pooled variance is:

$$\begin{aligned} s_p^2 &= \frac{(n_X - 1)s_X^2 + (n_Y - 1)s_Y^2}{n_X + n_Y - 2} = \frac{(21 - 1)1.30^2 + (25 - 1)1.16^2}{(21 - 1) + (25 - 1)} \\ &= 1.5021 \end{aligned}$$

- ▶ The test statistic is

$$t^{act} = \frac{(\bar{x} - \bar{y}) - d_0}{\sqrt{(s_p^2/n_X) + (s_p^2/n_Y)}} = \frac{(3.27 - 2.53) - 0}{\sqrt{1.5021 \left(\frac{1}{21} + \frac{1}{25}\right)}} = 2.040.$$

Since $|t^{act}| > t_{44,0.025} = 2.0154$, we reject H_0 at $\alpha = 0.05$. We conclude that there is evidence of a difference...

- The C.I. is constructed as $(\bar{X} - \bar{Y}) \pm t_{n_X+n_Y-2, \alpha/2} \cdot SE(\bar{X} - \bar{Y})$

Testing Population Mean Differences – I

Matched or Paired Samples

- Suppose we obtain a sample of n observations from two populations which are normally distributed and we have paired or matched samples – repeated measures (before/after).
- Define, the pair difference $d_i = X_i - Y_i$. We have

$$\bar{d} = \frac{1}{n} \sum_{i=1}^n d_i = \bar{X} - \bar{Y}; \quad \text{and} \quad S_d = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (d_i - \bar{d})^2}$$

with $E(\bar{d}) = \mu_d = E(X) - E(Y)$ and $SE(\bar{d}) = \sqrt{\frac{S_d^2}{n}} = S_d/\sqrt{n}$

- If the sample size is large enough ($n \rightarrow \infty$) then

$$\frac{\bar{d} - \mu_d}{S_d/\sqrt{n}} \sim N\left(0, \frac{S_d^2}{n}\right).$$

If the sample size is relatively small, then

$$\frac{\bar{d} - \mu_d}{S_d/\sqrt{n}} \sim t_{n-1}.$$

Testing Population Mean Differences – II


Matched or Paired Samples

$$\text{Matched or Paired Samples} \quad t = \frac{\bar{d} - d_0}{SE(d)} = \frac{\bar{d} - d_0}{s_d/\sqrt{n}} \quad (n \text{ large})$$

Lower-tail test:

$$H_0: E(X) - E(Y) \geq 0$$

$$H_1: E(X) - E(Y) < 0$$




Reject H_0 if $t < -z_\alpha$

Upper-tail test:

$$H_0: E(X) - E(Y) \leq 0$$

$$H_1: E(X) - E(Y) > 0$$




Reject H_0 if $t > z_\alpha$

Two-tail test:

$$H_0: E(X) - E(Y) = 0$$

$$H_1: E(X) - E(Y) \neq 0$$



Reject H_0 if $t < -z_{\alpha/2}$
or $t > z_{\alpha/2}$

Testing Population Mean Differences – III


Matched or Paired Samples

$$\text{Matched or Paired Samples} \quad t = \frac{\bar{d} - d_0}{\text{SE}(d)} = \frac{\bar{d} - d_0}{s_d/\sqrt{n}} \sim t_{n-1}$$

Lower-tail test:

$$H_0: E(X) - E(Y) \geq 0$$

$$H_1: E(X) - E(Y) < 0$$




$$-t_{n-1, \alpha}$$

Reject H_0 if $t < -t_{n-1, \alpha}$

Upper-tail test:

$$H_0: E(X) - E(Y) \leq 0$$

$$H_1: E(X) - E(Y) > 0$$




$$t_{n-1, \alpha}$$

Reject H_0 if $t > t_{n-1, \alpha}$

Two-tail test:

$$H_0: E(X) - E(Y) = 0$$

$$H_1: E(X) - E(Y) \neq 0$$



$$-t_{n-1, \alpha/2}$$

$$t_{n-1, \alpha/2}$$

Reject H_0 if $t < -t_{n-1, \alpha/2}$
or $t > t_{n-1, \alpha/2}$

Testing Population Mean Differences – I

Matched or Paired Samples: Example

- Assume you send your salespeople to a “customer service” training workshop. Has the training made a difference in the number of complaints? Test at the 5% significance level. You collect the following data:

Salesperson	C.B.	T.F	M.H.	R.K.	M.O.
Complaints, Before:	6	20	3	0	4
Complaints, After:	4	6	2	0	0
Difference, d_i	-2	-14	-1	0	-4

$$\bar{d} = \frac{1}{5} \sum_{i=1}^5 d_i = -4.2; \quad s_d = \sqrt{\frac{1}{5-1} \sum_{i=1}^5 (d_i - \bar{d})^2} = 5.67$$

- ▶ The hypothesis of interest is

$$H_0 : \mu_X - \mu_Y = 0$$

$$H_1 : \mu_X - \mu_Y \neq 0$$

Testing Population Mean Differences – II

Matched or Paired Samples: Example

- ▶ With $n = 4$ and $\alpha = 0.05$ the critical value is $t_{n-1, \alpha/2} = t_{4, 0.025} = 2.776$.
- ▶ We have

$$t = \frac{\bar{d} - d_0}{s_d / \sqrt{n}} = \frac{-4.2 - 0}{5.67 / \sqrt{4}} = -1.66 > -t_{4, 0.025} = -2.776,$$

or $|t| < t_{4, 0.025} = 2.776$. Hence, we **do not reject** H_0 . There is not a significant change in the number of complaints.

Annex: Hypothesis Tests – I

Employing the p -value

- Suppose we have a sample of n observations (they are assumed *IID*) and compute the sample average \bar{Y} . The sample average can differ from $\mu_{Y,0}$ for two reasons
 - 1 The population mean μ_Y is not equal to $\mu_{Y,0}$ (H_0 is not true)
 - 2 Due to random sampling $\bar{Y} \neq \mu_Y = \mu_{Y,0}$ (H_0 is true)
- To quantify the second reason we define the p -value. The *p -value* is the probability of drawing a sample with \bar{Y} at least as far from $\mu_{Y,0}$ as the value actually observed, given that the null hypothesis is true.

$$p\text{-value} = \Pr_{H_0} [|\bar{Y} - \mu_{Y,0}| > |\bar{Y}^{act} - \mu_{Y,0}|],$$

where \bar{Y}^{act} is the value of \bar{Y} actually observed

Annex: Hypothesis Tests – II

Employing the p -value

- To compute the p -value, you need to know the sampling distribution of \bar{Y} , which is complicated if n is small. With large n the CLT states that

$$\bar{Y} \sim N\left(\mu_Y, \frac{\sigma_Y^2}{n}\right),$$

which implies that if the null hypothesis is true:


$$\frac{\bar{Y} - \mu_{Y,0}}{\sqrt{\frac{\sigma_Y^2}{n}}} \sim N(0, 1)$$

- Hence

$$p\text{-value} = \Pr_{H_0} \left[\left| \frac{\bar{Y} - \mu_{Y,0}}{\sqrt{\frac{\sigma_Y^2}{n}}} \right| > \left| \frac{\bar{Y}^{act} - \mu_{Y,0}}{\sqrt{\frac{\sigma_Y^2}{n}}} \right| \right] = 2\Phi \left(- \left| \frac{\bar{Y}^{act} - \mu_{Y,0}}{\sqrt{\frac{\sigma_Y^2}{n}}} \right| \right)$$

Annex: Hypothesis Tests – III

Employing the p -value



- For large n , p -value = the probability that a $N(0, 1)$ random variable falls outside $\left| \frac{\bar{Y}^{act} - \mu_{Y,0}}{\sigma_{\bar{Y}}} \right|$, where $\sigma_{\bar{Y}} = \sigma_Y / \sqrt{n}$

Annex: Hypothesis Tests – I

Computing the p -value when σ_Y^2 is unknown

- In practice σ_Y^2 is usually unknown and must be estimated
- The sample variance S_Y^2 is the estimator of $\sigma_Y^2 = E[(Y - \mu_Y)^2]$, defined as

$$S_Y^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i - \bar{Y})^2$$

- ▶ division by $n - 1$ because we ‘replace’ μ_Y by \bar{Y} which uses up 1 degree of freedom
- ▶ if Y_1, Y_2, \dots, Y_n are *IID* and $E(Y^4) < \infty$, then $S_Y^2 \xrightarrow{p} \sigma_Y^2$ (Law of Large Numbers)
- The sample standard deviation $S_Y = \sqrt{S_Y^2}$, is the estimator of σ_Y .

Annex: Hypothesis Tests – II

Computing the p -value when $\sigma_{\bar{Y}}^2$ is unknown

- The standard error $SE(\bar{Y})$ is an estimator of $\sigma_{\bar{Y}}$

$$SE(\bar{Y}) = \frac{S_Y}{\sqrt{n}}$$

- Because S_Y^2 is a consistent estimator of σ_Y^2 we can (for large n) replace

$$\sqrt{\frac{\sigma_Y^2}{n}} \text{ by } SE(\bar{Y}) = \frac{S_Y}{\sqrt{n}}$$

- This implies that when σ_Y^2 is unknown and Y_1, Y_2, \dots, Y_n are *IID* the p -value is computed as

$$p\text{-value} = 2\Phi\left(-\left|\frac{\bar{Y}^{act} - \mu_{Y,0}}{SE(\bar{Y})}\right|\right)$$