

MSc Course: Mathematical Analysis - Optional Exercises (2020)

Stelios Arvanitis

- The following set of exercises is optional.
- The submission process will be announced.
- The instructor retains the right to ask for clarifications on the proposed solutions.

Exercise 1. Given an X -valued sequence $(x_n)_{n \in \mathbb{N}}$, define a $(x_n)_{n \in \mathbb{N}}$ -subsequence to be any infinite subset of $(x_n)_{n \in \mathbb{N}}$. Prove that if a metric space is totally bounded then every sequence has a Cauchy subsequence.

Exercise 2. Consider $(B(\mathbb{N}, \mathbb{R}), d_{\text{sup}})$ and $A = \left\{ (x_n)_{n \in \mathbb{N}}, x_n = \begin{cases} 1, & n = i \\ 2, & n \neq i \end{cases}, i \in \mathbb{N} \right\} \subset B(\mathbb{N}, \mathbb{R})$. Show that A is d_{sup} -bounded but not d_{sup} -totally bounded.

Exercise 3. Suppose that (X, d) is compact. Show that for $f_n : X \rightarrow \mathbb{R}$, $n \in \mathbb{N}$, if the sequence (f_n) is d_u/d -equi-Lipschitz then it is d_{sup} -bounded. Prove that if furthermore $f_n(x) \rightarrow f(x)$ as $n \rightarrow \infty$, for all $x \in X$, for some $f : X \rightarrow \mathbb{R}$, then also $d_{\text{sup}}(f_n, f) \rightarrow 0$ as $n \rightarrow \infty$, and conclude that the limit f is d_u/d -Lipschitz.

Exercise 4. Prove the Matkowski Fixed Point Theorem:

Theorem. Suppose that (X, d) is complete, $f : X \rightarrow X$, and $g : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that:

1. g is non-decreasing,
2. g is continuous at zero,
3. $g(t) = 0$ iff $t = 0$,
4. $\lim_{m \rightarrow \infty} g^{(m)}(t) = 0, \forall t \in \mathbb{R}_+$, and
5. $\forall t > 0, \lim_{m \rightarrow \infty} \frac{g^{(m+1)}(t)}{g^{(m)}(t)} = c_t < 1$.

Then if $\forall x, y \in X, d(f(x), f(y)) \leq g(d(x, y))$, f has a unique fixed point, say $x^* = \lim_{m \rightarrow \infty} f^{(m)}(x)$, for all $x \in X$.

Show that this is a generalization of BFPT.

Exercise 5. (Fredholm Integral Equation of the second kind.) Consider $X = C([a, b], \mathbb{R})$ with $d = d_{\text{sup}}$. Suppose that $\omega : [a, b] \times [a, b] \rightarrow \mathbb{R}$ is continuous, that $\omega(x, y) \geq 0$, $\forall x, y \in [a, b]$, that $0 < M_\omega := \sup_{x, y \in [a, b]} \omega(x, y)$, $h \in X$ and let $\lambda > 0$. Consider the integral equation

$$f(x) = h(x) + \lambda \int_a^b \omega(x, y) f(y) dy, \quad \forall x \in [a, b]. \quad (1)$$

Show that there exists a unique $f \in X$ that satisfies (1) if $\lambda < \frac{1}{M_\omega(b-a)}$.