

Problem Set 1**

Metric functions and metric spaces

Exercise 1

Is $d(x, y) = |x - y|$ a metric?

Exercise 2

Is the function $d : X \times X \rightarrow \mathbb{R}$ such that $d(x, y) = |x - y|, \forall x, y \in X$ a metric on the non-empty set $X \subseteq \mathbb{R}$?

Exercise 3

Suppose that (Y, d) is a metric space. Let $f : X \rightarrow Y$ be an injection from X to Y . Define $d_f : X \times X \rightarrow \mathbb{R}$ such that $d_f(x, y) = d(f(x), f(y)), \forall x, y \in X$. Is (X, d_f) a metric space?

Exercise 4

Study whether or not the following pairs of sets and functions constitute metric spaces:

1. $X \neq \emptyset$ and $d(x, y) = \begin{cases} 0, & x = y \\ c, & x \neq y \end{cases}, \forall x, y \in X$, with $c > 0$ (Discrete distance)
2. $X = \mathbb{R}$ and $d(x, y) = |e^x - e^y|, \forall x, y \in X$ [Sutherland Ex. 5.4 (b)]
3. $X = \emptyset$ and $d(x, y) = |x - y|, \forall x, y \in X$
4. $X = \mathbb{R}$ and $d(x, y) = \ln(|e^x - e^y|), \forall x, y \in X$
5. $X = [-1, 1]$ and $d(x, y) = |x^2 - y^2|, \forall x, y \in X$
6. $X = \mathbb{R}$ and $d(x, y) = |x - y^3|, \forall x, y \in X$

*Please report any typos, mistakes, or even suggestions at zaverdasd@aueb.gr.

**Some exercises were collected and compiled by Dr. Alexandros Papadopoulos.

7. $X = [0, 1]$ and $d(x, y) = |x - y|^2, \forall x, y \in X$

8. $X = \mathbb{R}^N$ and $d(x, y) = \left(\sum_{i=1}^N |x_i - y_i|^p\right)^{\frac{1}{p}}, \forall x, y \in X$, with $p, N \in \mathbb{N}^*$ (Minkowski distance)

Exercise 5

For any metric space (X, d) and $\forall x, y, z, w \in X$, show that:

1. $|d(x, z) - d(z, y)| \leq d(x, y)$ [O'Searcoid Theorem 1.1.2, Sutherland Ex. 5.1]
2. $|d(x, y) - d(z, w)| \leq d(x, z) + d(y, w)$ [O'Searcoid Q 1.2, Sutherland Ex. 5.2]

Exercise 6

Let X be some non-empty set. Let d_1, d_2 , and d_s be distance functions on X such that $d_s = d_1 + d_2$. Determine whether the following statements always hold (or under which conditions they could hold):

1. If d_1 and d_2 are metrics on X , d_s is a metric on X .
2. If d_1 is a metric and d_2 a pseudo-metric on X , d_s is a metric on X .
3. If d_1 and d_2 are pseudo-metrics on X , d_s is a metric on X .

Exercise 7

Consider a finite index set $\mathcal{I} = \{1, 2, \dots, n\}$ with $n \in \mathbb{N}^*$ and for each of its elements, i , the functional metric spaces $(\mathcal{B}(X_i, \mathbb{R}), d_{sup}^i)$ with

$$d_{sup}^i(f_i, g_i) = \sup_{x \in X_i} |f_i(x) - g_i(x)|, \forall f_i, g_i \in \mathcal{B}(X_i, \mathbb{R})$$

Consider the product set $B_{\Pi} := \prod_{i \in \mathcal{I}} \mathcal{B}(X_i, \mathbb{R})$ with $f := (f_i)_{i \in \mathcal{I}} \in B_{\Pi}$ and the function $d_{\Pi} : B_{\Pi} \times B_{\Pi} \rightarrow \mathbb{R}$ such that

$$d_{\Pi}(f, g) = \max_{i \in \mathcal{I}} \sup_{x \in X_i} |f_i(x) - g_i(x)|, \forall f, g \in B_{\Pi}$$

Is (B_{Π}, d_{Π}) a metric space?

Exercise 8 [O'Searcoid Q 1.8]

Let $P(S)$ be the power set of a non empty set, S . Let the function $d : P(S) \times P(S) \rightarrow \mathbb{R}$ such that

$$d(A, B) = |(A \setminus B) \cup (B \setminus A)|, \forall A, B \in P(S)$$

be a function that gives the cardinality of the symmetric difference between two elements of $P(S)$ (i.e. subsets of S). Is d a metric on $P(S)$?

Exercise 9 [Sutherland Ex. 5.14]

Let n be a positive natural number. The distance functions:

1. $d_1 : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ such that $d_1(x, y) = \sum_{i=1}^n |x_i - y_i|$, $\forall x, y \in \mathbb{R}^n$ (Manhattan distance)
2. $d_2 : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ such that $d_2(x, y) = \sqrt{\sum_{i=1}^n |x_i - y_i|^2}$, $\forall x, y \in \mathbb{R}^n$ (Euclidean distance)
3. $d_\infty : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ such that $d_\infty(x, y) = \max_{i=1}^n |x_i - y_i|$, $\forall x, y \in \mathbb{R}^n$ (Chebyshev distance)

are all metrics on \mathbb{R}^n . Show that the following functional inequalities hold:

$$d_\infty \leq d_2 \leq d_1 \leq n \cdot d_\infty \leq n \cdot d_2 \leq n \cdot d_1$$

Exercise 10

Let X be an $n \times m$ real matrix, with $n, m \in \mathbb{N}^*$ and $n > m$, such that $\text{rank}(X) = m$. Then $P_X = X(X'X)^{-1}X'$ is the projection matrix of X . Let $Y \subseteq \mathbb{R}^n$ be non-empty and \hat{Y} be its projected image through P_X . Define $d_X : Y \times Y \rightarrow \mathbb{R}$ such that $d_X(x, y) = \|P_X \cdot x - P_X \cdot y\|$, $\forall x, y \in Y$ (i.e. d_X is the Euclidean norm of an n -dimensional real vector). Show that (Y, d_X) is a pseudo-metric space.

(Hint: Consider the example of exercise 3. Under which conditions for f is (X, d_f) a pseudo-metric space?)

Exercise 11

Let (X, d) be a metric space and consider a real function $f : \mathbb{R} \rightarrow \mathbb{R}$. Define $d' : X \times X \rightarrow \mathbb{R}$ such that $d'(x, y) = f(d(x, y))$, $\forall x, y \in X$.

1. Deduce the necessary conditions for f for d' to be a metric on X .
2. Is it a sufficient condition for f to be a strictly increasing concave real function with $f(0) = 0$ for d' to be a metric on X ?

Useful Theorems and Results

Cardinality and Set Operations

Cardinality is a measure of the number of elements in a set. The following properties hold with respect to cardinality:

$$|\emptyset| = 0 \tag{1}$$

$$|A| + |B| = |A \cup B| + |A \cap B| \tag{2}$$

$$|A \setminus B| = |A| - |A \cap B| \tag{3}$$

Square of the sum of N numbers

$$\left(\sum_{i=1}^N a_i \right)^2 = \sum_{i=1}^N a_i^2 + 2 \sum_{i=1}^N \sum_{j=1}^{i-1} a_i a_j \tag{4}$$

Hölder's inequality

For all $x, y \in \mathbb{R}^N$ and $\alpha, \beta \in (1, +\infty)$ such that $\frac{1}{\alpha} + \frac{1}{\beta} = 1$, it holds that

$$\sum_{i=1}^N |x_i y_i| \leq \left(\sum_{i=1}^N |x_i|^\alpha \right)^{\frac{1}{\alpha}} \left(\sum_{i=1}^N |y_i|^\beta \right)^{\frac{1}{\beta}} \tag{5}$$

For $\alpha = \beta = 2$ we get the Cauchy-Schwartz inequality.

For $x \in \mathbb{R}^N$ we call $\|x\|_p := \left(\sum_{i=1}^N |x_i|^p \right)^{\frac{1}{p}}$ the p -norm of x .